

LA-UR-21-20608

Approved for public release; distribution is unlimited.

Title: Introduction to FEL Simulations

Author(s): Anisimov, Petr Mikhaylovich
Neveu, Nicole
Nguyen, Dinh C.

Intended for: USPAS 21 lecture notes

Issued: 2021-01-25

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



VUV and X-ray Free-Electron Lasers

Introduction to FEL Simulations

Petr Anisimov,² Nicole Neveu¹, Dinh C. Nguyen¹

¹ SLAC National Accelerator Laboratory

² Los Alamos National Laboratory



Wednesday (Jan 27) Lecture Outline

- Canonical formulation of 1D FEL theory
- Q&A
- Numerical simulator for 1D FEL
- Q&A
- Numerical simulator – ZFEL code
- Q&A

Canonical formulation of 1D FEL theory

Lagrangian description

- Lagrangian for a relativistic electron in an electromagnetic field:^{1,2}

$$L(\mathbf{r}, \dot{\mathbf{r}}; t) = -m_e c^2 \sqrt{1 - \frac{\dot{\mathbf{r}}^2}{c^2}} + e\phi(\mathbf{r}) - e\dot{\mathbf{r}} \cdot \mathbf{A}(\mathbf{r})$$

- Lagrangian density for classical electrodynamics:^{1,3}

$$\mathcal{L} = -\frac{1}{4\mu_0} F^{\alpha\beta} F_{\alpha\beta} - A_\alpha J^\alpha,$$

where, $F^{\alpha\beta} F_{\alpha\beta} = 2(\partial_\alpha A_\beta \partial^\alpha A^\beta - \partial_\beta A_\alpha \partial^\alpha A^\beta)$, $A_\alpha = \left(\frac{\phi(\mathbf{r})}{c}, -\mathbf{A}(\mathbf{r})\right)$ and $J^\alpha = (c\rho(\mathbf{r}), \mathbf{j}(\mathbf{r}))$;

- The Hamiltonian principle: $\delta S = \delta \int_{t_1}^{t_2} L(\mathbf{r}, \dot{\mathbf{r}}; t) dt = 0$.

¹ L. D. Landau and E. M. Lifshitz, *The classical theory of fields*; 3rd ed., ser. Course of theoretical physics. Oxford: Pergamon, 1971, Eq. (28.6) [Conversion of Gaussian to SI units is discussed at https://en.wikipedia.org/wiki/Gaussian_units#Major_differences_between_Gaussian_and_SI_units]

² https://en.wikipedia.org/wiki/Relativistic_Lagrangian_mechanics and ³ https://en.wikipedia.org/wiki/Covariant_formulation_of_classical_electromagnetism

FEL in Lagrangian description

- An ensemble of electrons creates charge density $\rho(\mathbf{r}) = -e \sum_{n=1}^{N_e} \delta(\mathbf{r} - \mathbf{r}_n(t))$ and the current density $\mathbf{j}(\mathbf{r}) = -e \sum_{n=1}^{N_e} \dot{\mathbf{r}}_n(t) \delta(\mathbf{r} - \mathbf{r}_n(t))$ resulting in

$$L_{int} = - \int A_\alpha J^\alpha d^3\mathbf{r} = e \sum_{n=1}^{N_e} [\phi(\mathbf{r}_n) - \dot{\mathbf{r}}_n \cdot \mathbf{A}(\mathbf{r}_n)],$$

which is the same term as in the Lagrangian for a relativistic electron in an electromagnetic field but summed over all the particles;

- There are no stationary charges therefore $\phi(\mathbf{r}) = 0$ will be assumed resulting in Coulomb and Lorentz gauges¹ becoming $\nabla \cdot \mathbf{A} = 0$.

¹ https://en.wikipedia.org/wiki/Gauge_fixing

² For more info on the Electromagnetic tensor $F^{\alpha\beta}$ go to https://en.wikipedia.org/wiki/Electromagnetic_tensor

Assumptions for 1D FEL

- Consider a helical undulator with a period λ_u that is described by

$$\mathbf{A}_u(z) = \frac{m_e c}{\sqrt{2} e} K \hat{\epsilon} e^{-ik_u z} + c.c.,$$

where $\hat{\epsilon} = (\hat{x} + i\hat{y})/\sqrt{2}$ is the polarization vector, $k_u = 2\pi/\lambda_u$ is the undulator wavenumber and $K = 0.934 B_0[T]\lambda_u[cm]$ is the undulator parameter;

- Further consider a plane-wave radiation field of wavelength λ :

$$\mathbf{A}_r(z, t) = -\frac{i}{\sqrt{2} k} E(t) \hat{\epsilon} e^{ikz - i\omega t} + c.c.,$$

where the phase convention is such that $\mathbf{A}_u \cdot \mathbf{A}_r \neq 0$.

- Since the Lagrangian does not depend on x or y variables then the canonical momenta $\frac{\partial L}{\partial \dot{\mathbf{r}}_\perp} \stackrel{\text{def}}{=} \mathbf{p}_\perp = \text{const}$ and could be set to zero.

1D FEL in Lagrangian description

- The undulator vector potential is externally created and does not have to be included in the Lagrangian of the system. The Lagrangian of the radiation field confined to some volume V is

$$L = \int \mathcal{L} d\mathbf{r} = -\frac{1}{2\mu_0} \left(B_r^2 - \frac{E_r^2}{c^2} \right) V \approx \frac{V}{\mu_0} \frac{iE^*}{\omega} \dot{E},$$

which results in the canonical momentum for the electromagnetic field to be

$$p_E \stackrel{\text{def}}{=} \frac{dL}{d\dot{E}} = \frac{V}{\mu_0} \frac{iE^*}{\omega};$$

- The Hamiltonian principle can now be rewritten as

$$\delta \left(\int_{t_1}^{t_2} p_E dE + \sum_{n=1}^{N_e} p_{z,n} dz_n - H_n dt \right) = 0.$$

1D FEL in Hamiltonian description

- The Hamiltonian for a single electron becomes

$$H_n \approx c \sqrt{m^2 c^2 + p_{z,n}^2 + 2 \frac{e m_e K}{k} \text{Im}[E(t) e^{i\theta_n}]},$$

where $\theta_n = (k + k_u)z_n - \omega t$ is the ponderomotive phase; $m^2 = m_e^2(1 + K^2)$ is a longitudinal mass of an ‘undulator’ electron and we have neglected $\sim |E(t)|^2$ term;

- We will replace the canonical variable z_n with θ_n in the Hamiltonian principle such that $dz_n = (d\theta_n + \omega dt)/(k + k_u)$;
- We will define the efficiency of an FEL interaction $\rho = \frac{E_{sat}^2 V}{\mu_0} / \gamma_0 m_e c^2 N_e$ as the ratio of the EM energy at saturation to the beam energy and rescale the field amplitude to its value at saturation $dE = E_{sat} dA$.

1D FEL in Hamiltonian description cont'd

- The Hamiltonian principle in the FEL theory notations becomes

$$\delta \left(\int_{\tau_1}^{\tau_2} p_A dA + \sum_{n=1}^{N_e} p_{\theta,n} d\theta_n - \tilde{H}_n d\tau \right) = 0$$

in terms of scaled time $dt = d\tau / 2\rho k_u c$, new canonical momenta $p_{\theta,n} = \frac{p_{z,n}}{k+k_u}$ and $p_A = i \frac{\rho \gamma_0 m_e c}{k} N_e A^*$, and a new Hamiltonian for n^{th} electron


$$\tilde{H}_n = \frac{1}{2k_u \rho} \left(\sqrt{(k+k_u)^2 p_{\theta,n}^2 + m^2 c^2 (1 + \mathcal{V})} - k p_{\theta,n} \right),$$

where a scaled ponderomotive potential has been introduced

$$\mathcal{V} = \frac{2K}{1+K^2} \frac{\omega_{pe}}{\omega} \sqrt{\rho \gamma_0} \text{Im}[A(\tau) e^{i\theta_n}],$$

in terms of the beam plasma frequency $\omega_{pe}^2 = \frac{e^2}{m_e \epsilon_0} \frac{N}{V}$.

FEL Hamiltonian



- In the absence of the ponderomotive potential created by the radiation the Hamiltonian of n^{th} electron becomes:

$$\tilde{H} = \frac{mc}{2k_u\rho} \left(\sqrt{1 + p^2} - v_{BR}p \right),$$

where a new momentum $p = \frac{(k+k_u)p_\theta}{mc}$ and the Bambini-Renieri velocity in units of speed of light $v_{BR} = \frac{k}{k+k_u}$ have been introduced;

- We here illustrate the case of $\lambda = 0.3 \text{ A}$ radiation in $\lambda_u = 1.86 \text{ cm}$ undulator.

Ponderomotive Potential

- The scaled ponderomotive potential $\mathcal{V} = \frac{2K}{1+K^2} \frac{\omega_{pe}}{\omega} \sqrt{\rho\gamma_0} \text{Im}[A(\tau)e^{i\theta_n}] \ll 1$ can be treated as a perturbation to the Hamiltonian evolution:

$$\frac{dp_\theta}{d\tau} = -\frac{\partial \tilde{H}}{\partial \theta} = 0$$

$$\frac{d\theta}{d\tau} = \frac{\partial \tilde{H}}{\partial p_\theta} = \frac{k + k_u}{2k_u\rho} \left(\frac{p}{\sqrt{1+p^2}} - v_{BR} \right)$$

where the canonical momentum is conserved and the phase is linearly increasing;

- Efficient interaction with the Ponderomotive potential requires $\langle \mathcal{V} \rangle_\theta \neq 0$, which could be achieved with $\frac{d\theta}{d\tau} = 0 = \frac{\partial \tilde{H}}{\partial p_\theta}$ condition that corresponds to the minimum of \tilde{H} !

The resonant energy and the FEL parameter

- The equilibrium momentum that minimizes the Hamiltonian and maximizes the ponderomotive interaction can be expressed as $p_{\theta}^{eq} = \frac{\gamma_{eq} m \dot{z}_{eq}}{k + k_u}$ with an equilibrium Lorentz factor γ_{eq} corresponding to the velocity $\dot{z}_{eq} = c v_{BR}$;
- This implies that the resonant energy for an electron is $\gamma_r^2 \approx (1 + K^2)k/2k_u$ in accordance with the classical X-ray FEL theory!
- The final Hamiltonian thus becomes

$$H_n = H_0 + \frac{p_n^2}{2M} + 2M \operatorname{Im}[A(\tau)e^{i\theta_n}],$$

where $H_0 = mc^2/2k_u c \rho \gamma_{eq}$, p_n is the detuning for the equilibrium momentum,

and $M = \rho p_{\theta}^{eq}$ if one chooses the FEL parameter to be $\rho = \frac{1}{\gamma_r} \left(\frac{K \omega_{pe}}{4ck_u} \right)^{\frac{2}{3}}!$

1D FEL equations

$$\begin{aligned} \frac{dA}{d\tau} &= \frac{\partial}{\partial p_A} \sum_{n=1}^{N_e} H_n \approx \frac{1}{N_e} \sum_{n=1}^{N_e} e^{-i\theta_n} \\ \frac{d\theta_n}{d\tau} &= \frac{\partial H_n}{\partial p_n} = \frac{p_n}{M} \\ \frac{dp_n}{d\tau} &= -\frac{\partial H_n}{\partial \theta_n} = -2M \operatorname{Re}[A(\tau)e^{i\theta_n}] \end{aligned}$$

One can introduce a scaled energy detuning $\eta_n = \frac{p_n}{M} = \frac{\gamma_n - \gamma_r}{\rho \gamma_r}$ in order to recover the Eqs. 4.31a, 4.31b, 4.31c and 4.31d of the book.

Additional reading on The FEL Hamiltonian

- S. D. Webb, “Period-Averaged Symplectic Maps for the FEL Hamiltonian” 38th International Free Electron Conference, 2017;
- P. M. Anisimov, “Canonical Formulation of 1D FEL Theory Revisited, Quantized and Applied to Electron Evolution”, 38th International Free Electron Conference, 2017;
- P. M. Anisimov, “Quantum theory for 1D X-ray Free Electron Laser”, Journal of Modern Optics, 65(11), pp 1370-1377, 2018.

Numerical simulator for 1D FEL

1D FEL equations in Python

$$\frac{dA}{d\tau} = \frac{1}{N_e} \sum_{n=1}^{N_e} e^{-i\theta_n}$$

$$\frac{d\theta_n}{d\tau} = \eta_n$$

$$\frac{d\eta_n}{d\tau} = -2 \operatorname{Re}[A(\tau)e^{i\theta_n}]$$

```
from scipy.integrate import solve_ivp
import numpy as np
import matplotlib.pyplot as plt

def rhs(t, y):
    """
    The right-hand side of the 1D canonical FEL equations;
    t - the current time;
    y - array of [A, theta, eta]
    """
    n = len(y)//2
    A = y[0]
    theta = y[1:n+1]
    eta = y[n+1:]
    dA_dt = np.mean(np.exp(-1j*theta))
    dtheta_dt = eta
    deta_dt = -2*np.real(A*np.exp(1j*theta))
    return np.concatenate(( [dA_dt],
                             dtheta_dt,
                             deta_dt))



sol = solve_ivp(rhs, [0, 4*np.pi],
                 np.concatenate(( [A0+0j], theta0, eta0)),
                 max_step=0.1)
```

What are the initial conditions?

- $A(0) = A_0$ corresponds to a seeded FEL and $A(0) = 0$ corresponds to SASE FEL;
- $\theta_n = 0$ gives bunching $b = \frac{1}{N_e} \sum_{n=1}^{N_e} e^{-i\theta_n} = 1$;
- A random uniform distribution $\theta_n \in [-\pi, \pi)$ has bunching $b = \frac{1}{N_e} \sum_{n=1}^{N_e} e^{-i\theta_n} = 0$ but $\langle |b|^2 \rangle_\theta = 1/N_e$, which is sufficient to initiate FEL instability;
- $\theta_n = x_n + \delta\theta \sin(x_n)$ such that $\langle |b|^2 \rangle = 0$ when $\delta\theta = 0$ – a so called ‘quiet start’.
- What other initial conditions can you think of?


Case 1

```
ne = 100
tau = 20
A0 = 0
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```




Case 1

```
ne = 100
tau = 20
A0 = 0
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```




Case 2 – “quiet start”

```
ne = 100
tau = 20
A0 = 0
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne, endpoint=False)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```




Case 3 – “seeded operation”

```
ne = 100
tau = 20
A0 = 1e-3
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne, endpoint=False)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```




Case 3 – “seeded operation”

```
ne = 100
tau = 20
A0 = 1e-3
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne, endpoint=False)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```



Case 3 – “seeded operation” scan



Observe:

- Lethargy period;
- Exponential growth;
- Saturated power;
- Power circulation.

Compare:


$$P = \frac{P_0}{9} \left| e^{\frac{i+\sqrt{3}}{2}\tau} + e^{\frac{i-\sqrt{3}}{2}\tau} + e^{-i\tau} \right|^2$$

Invariant of the evolution:

$$\eta(\tau) + |A(\tau)|^2 = \text{const}$$


Case 4 – “SASE operation”

```
ne = 100
tau = 5*np.pi
A0 = 0
theta0 = np.random.uniform(low=-4*np.pi, high=4*np.pi, size=ne)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```




Case 4 – “SASE operation”

```
ne = 100
tau = 5*np.pi
A0 = 0
theta0 = np.random.uniform(low=-4*np.pi, high=4*np.pi, size=ne)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```




Case 4 – “SASE operation” scan

```
ne = 100
tau = 5*np.pi
A0 = 0
theta0 = np.random.uniform(low=-4*np.pi, high=4*np.pi, size=ne)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```





Case 5 – “Energy spread driven FEL” scan

```
ne = 100
tau = 5*np.pi
A0 = 0
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne, endpoint=False)
p0 = np.random.normal(0, 0.1, ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```



Case 6 – “controlled SASE”

```
ne = 100
tau = 5*np.pi
A0 = 0
dtheta = 2/np.sqrt(100)
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne, endpoint=False)
theta0 -= dtheta*np.sin(theta0)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
```



Case 6 – “controlled SASE”



```
ne = 100
tau = 5*np.pi
A0 = 0
dtheta = 2/np.sqrt(100)
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne, endpoint=False)
theta0 -= dtheta*np.sin(theta0)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
plot(sol)
```

- The choice of $\delta\theta = 2/\sqrt{N_e}$ comes from the requirement to recover the correct Poisson statistics¹ such that $|\langle e^{-i\theta_n} \rangle| = \frac{1}{2}\delta\theta = 1/\sqrt{N_e}$;
- N_e is the number of interacting electrons such that $N_e = 4.3 \frac{L_g}{\lambda_u} \frac{I_b \lambda}{e v_b}$ and $\delta\theta = 1.38 \times 10^{-3}$ corresponds to MaRIE XFEL case;
- We can also define an equivalent startup noise power $|A(0)|^2 = N_e^{-1}$.

¹We refer an interested reader to “The effect of shot noise on the start up of the fundamental and harmonics in free-electron lasers” by H. P. Freund et al., J. Appl. Phys. 104, 123114 (2008); <https://doi.org/10.1063/1.3040689>


Case 7 – SASE vs shot noise power

```
ne = 25
tau = 5*np.pi
A0 = 0
dtheta = 1.38e-3
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne, endpoint=False)
theta0 -= dtheta*np.sin(theta0)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
```



Case 7 – SASE vs shot noise power


```
ne = 25
tau = 5*np.pi
A0 = 0
dtheta = 1.38e-3
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne, endpoint=False)
theta0 -= dtheta*np.sin(theta0)
p0 = np.zeros(ne)
sol = solve_ivp(rhs, [0, tau], np.concatenate(([A0+0j], theta0, p0)), max_step=0.1)
```




Observe:

- SASE vs shot noise power distinction happens during the lethargy period;
- The number of macro particles used for the simulation does not depend on the number of interacting electrons $n_e \ll N_e$!

Case 7 – SASE vs shot noise power




Numerical simulation vs Analytical solution




Phase in SASE case (green) vs Seeded case (blue)

Case 7 – A typical animation




Case 8 – resonance curve

```
ne = 25
tau = 10*np.pi
A0 = 1.38e-3/2
dtheta = 0
theta0 = np.linspace(-4*np.pi, 4*np.pi, ne, endpoint=False)
theta0 -= dtheta*np.sin(theta0)
res = []
eta = np.linspace(-2, 2.3)
for eta0 in eta:
    p0 = eta0*np.ones(ne)
    sol = solve_ivp(rhs, [0, tau],
                    np.concatenate([[A0+0j], theta0, p0),
                    max_step=0.1)
    res.append(np.max(np.abs(sol.y[0])**2))
plt.plot(eta, res, 'o-')
plt.xlabel(r'Energy detuning, $\eta_0$')
plt.ylabel(r'Peak power, $\max(|A|^2)$')
plt.xlim([-2, 2.3])
plt.ylim([0, 4])
plt.show()
```



Case 9 – correct resonance curve (HW)



Observe:

- Maximum gain of $\sqrt{3}$ is reached at zero energy detuning;
- The resonance curve is asymmetric;
- The lower energy case keeps lasing longer!

Numerical simulator – ZFEL code

<https://github.com/slaclab/zfel.git>


ZFEL package structure

- `sase1d_input_part.sase` performs 1D FEL simulation
 - `.sase.params_calc` performs FEL normalization
 - `.general_load_bucket.general_load_bucket` performs (θ_n, η_n) distribution generation
 - `.sase.FEL_process` performs dimensionless FEL calculations
 - `.sase.final_calc` converts dimensionless results back to physical units
- Output:
 - `z, power_z, s, power_s, rho, detune, field, field_s, gainLength, resWavelength, thet_out, eta_out, bunching, spectrum, freq, Ns, history`

SASE 1D FEL run function: Input



Electron beam description		X-ray radiation description	
energy	electron energy [eV]	s_steps	n-sample points along bunch length
eSpread	relative rms energy spread []	radWavelength	seed wavelength? [meter], used only in single-frequency runs
emitN	normalized transverse emittance [m-rad]	iopt	'sase' or 'seeded'
currentMax	peak current [Ampere]	P0	small seed input power [W]
beta	mean beta [meter]		
Undulator description		Technical description	
z_steps	n-sample points along undulator	Nruns	not implemented yet
unduPeriod	undulator period [meter]	npart	n-macro-particles per bucket
unduk	undulator parameter, array of K []	constseed	use constant random seed for reproducibility, 1 Yes, 0 No
undul	length of undulator [meter]	particle_position	particle distro with positions in meter and eta
dEdz	not implemented yet	hist_rule	different rules to select number of intervals to generate the histogram of eta value in a bucket

Graphical representation



Slippage

- The FEL resonance condition $(k + k_u)z - \omega_x t = \text{const}$ means that an x-ray radiation overtakes e-beam by $\delta s = \lambda_x$ every $\Delta z = \lambda_u$;
- Therefore, $\delta s = \Delta z \lambda_x / \lambda_u$ is the distance that the radiation overtakes the e-beam every Δz integration step.



ZFEL code setup

Undulator description

```

unduPeriod = 1.86e-2 # undulator period []
unduL = 3500*unduPeriod # undulator length [m]
z_steps = 100 # n-sample points along undulator; it also defines time step along bunch
unduK = np.full(z_steps, np.sqrt(2)*0.86) # undulator parameter
dEdz = None # not implemented

```

Electron beam description

```

currentMax = 3e3 # maximum current [A]
energy = 12e9 # electron energy [eV]
eSpread = 1.5e-4 # relative rms energy spread sigma_E/E
emitN = 0.2e-6 # normalized emittance [m-rad]
beta = 15 # mean beta function of the beam [m]

```

ZFEL code setup cont'd

X-ray radiation description

```
iopt = 'sase' # The only mode of operation
P0 = 0e4 # small seed input power [W]
radWavelength = 2.934e-11 # seed wavelength? [meter],
                        # used only in single-freugency runs
s_steps = 10000 # n-samples along bunch; tail is at 0; it also defines bunch length
```

Technical description


```
Nruns = None # not implemented
npart = 2**9 # n-macro-particles per bucket
constseed = 1 # use constant random seed for reproducibility,
              # 1 Yes, 0 No
particle_position = None # particle distro with positions in
                        # meter and eta
hist_rule = 'square-root' # 'square-root' or 'sturges' or
                        # 'rice-rule' or 'self-design',
                        # number of intervals to generate
                        # the histogram of eta value in a bucket
```


Execution

Execute



```
# Load input parameters into zfel
inp_struct={'Nruns':Nruns, 'npart':npart, 's_steps':s_steps, 'z_steps':z_steps, \
            'energy':energy, 'eSpread':eSpread, 'dEdz':dEdz, \
            'emitN':emitN, 'currentMax':currentMax, 'beta':beta, \
            'unduPeriod':unduPeriod, 'unduK':unduK, 'unduL':unduL, \
            'radWavelength':radWavelength, 'iopt':iopt, 'P0':P0, \
            'constseed':constseed, 'particle_position':particle_position, 'hist_rule':hist_rule}
z, power_z, s, power_s, rho, detune, field, \
field_s, gainLength, resWavelength, \
thet_out, eta_out, bunching, spectrum, freq, Ns, history = sase1d_input_part.sase(inp_struct)
```

Results: x-ray power




Simulated power at a MaRIE-like x-ray FEL in the 1D approximation. Strong SASE power fluctuations are present. Averaged power exhibits expected characteristics of initial lethargy, exponential growth and saturation.

Results: phase space distribution



Results: spectral analysis



Simulated power spectrum at a MaRIE-like x-ray FEL in the 1D approximation. Strong SASE power fluctuations corresponding to the spectrum with a relative bandwidth of 4×10^{-4} . The contrast of the resonant radiation vs incoherent background is about 10^{-4} .

Results: Transverse dynamics?

